

Table Yersiniosis notifications 1996-2005 and source of infection (definite and suspected)

Report year	Total no. notified cases (N)	Source identified as 'No Source' or 'Unknown'		Source identified as 'Definite' or 'Suspect'		Detail for source identified as 'Definite' or 'Suspect' *									
						Source: Person to Person		Source: Food or Drink		Source: Animal		Source: Other specified		Source: Not specified	
		No.	% of total	No.	% of total	No.	%	No.	%	No.	%	No.	%	No.	%
1996	330	298	90.3%	32	9.7%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	32	100.0%
1997	488	425	87.1%	63	12.9%	11	17.5%	24	38.1%	0	0.0%	0	0.0%	30	47.6%
1998	546	481	88.1%	65	11.9%	15	23.1%	52	80.0%	0	0.0%	0	0.0%	0	0.0%
1999	503	409	81.3%	94	18.7%	11	11.7%	69	73.4%	22	23.4%	8	8.5%	1	1.1%
2000	396	328	82.8%	68	17.2%	7	10.3%	47	69.1%	17	25.0%	8	11.8%	1	1.5%
2001	429	357	83.2%	72	16.8%	11	15.3%	41	56.9%	27	37.5%	14	19.4%	1	1.4%
2002	476	422	88.7%	54	11.3%	3	5.6%	25	46.3%	20	37.0%	13	24.1%	0	0.0%
2003	439	396	90.2%	43	9.8%	7	16.3%	20	46.5%	22	51.2%	6	14.0%	0	0.0%
2004	420	370	88.1%	50	11.9%	3	6.0%	21	42.0%	23	46.0%	6	12.0%	1	2.0%
2005	407	348	85.5%	59	14.5%	1	1.7%	25	42.4%	23	39.0%	14	23.7%	5	8.5%
Grand Total	4434	3834	86.5%	600	13.5%	69	11.5%	324	54.0%	154	25.7%	69	11.5%	71	11.8%

Source: EpiSurv data as at 9 May 2006, Institute of Environmental Science and Research Ltd.

Notes:

Annual figures may differ from published ESR reports as notification data can be updated or changed by a public health service if further information becomes available.

* Total across the sources in this section may be greater than the total in the Source identified as 'Definite' or 'Suspect' section due to multiple sources for a single case being reported.